

National Cervical Screening Management Guide



National Cervical Screening Guidelines

A Brief Overview

In December 2017 the Pap smear process was replaced with the Cervical Screening Test. The cervical cells are collected in the same way, with the only difference being that the sample is placed in a ThinPrep® vial with liquid media rather than smeared onto glass slides. Testing the sample for oncogenic Human papillomavirus (HPV) replaced cytology as the primary screening process. Liquid-based cytology (LBC) is performed on the sample by the laboratory if the HPV test is positive (i.e. reflex cytology) or in specific clinical circumstances (i.e. HPV and cytology co-test).

The cervical screening program is available to women between the ages of 25 and 74 years. Women over the age of 25 are invited by the National Cancer Screening Register to participate in the National Cervical Screening Program. The recommended interval between cervical screening tests is 5 years.*

Women who have had the HPV vaccination must still participate in the screening program.

A detailed explanation of the program is available from Cancer Council Australia.

HPV testing

Human papillomavirus (HPV) is a common virus. Most infections are harmless and resolve spontaneously in about a year. In some patients, persistent infection with one of the oncogenic genotypes of HPV can lead to cervical pre-cancer or cancer. Types 16 and 18 are more virulent than other HPV types, consistently causing around 75% of all cervical cancers.*

Testing for oncogenic HPV types has been shown to be as sensitive as cytology in identifying women at risk of developing cervical neoplasia. However, it is the strong negative predictive value of HPV testing that has the most clinical use.*

QML Pathology use the latest HPV testing technology for the detection of the 14 HPV genotypes known to be associated with cervical cancer. The test specifically detects HPV 16 and 18 (which cause 75% of cervical cancers) while simultaneously detecting the 12 other oncogenic genotypes. There is an internal control which minimises the risk of false negative results for each patient.

Remember, a positive HPV result does not necessarily indicate that the woman has cervical neoplasia, but does indicate an increased risk.

Routine 5-yearly Cervical Screening

For women aged 25 to 69

Depending on the result of the HPV, and any additional reflex cytology testing, women are assigned a clinical risk category – Low Risk, Intermediate Risk or Higher Risk. Each risk category follows a different clinical pathway:

Cervical Screening Pathways

1. Low Risk = the HPV test is negative → The woman will be invited to rescreen in five years

 **Low Risk**

2. Intermediate Risk = the HPV test is positive for one of the other oncogenic HPV types (i.e. NOT 16/18) and the reflex cytology performed by the lab is either negative or only shows low grade changes.

 **Intermediate Risk**


- The woman will be invited to have another HPV test in 12 months
- If the repeat HPV test in 12 months is negative then the woman can return to routine 5 yearly screening
- If the repeat HPV test in a 12 months shows the same criteria for intermediate risk as above, then they are invited to have another HPV test in 12 months
- If that second repeat HPV test in 12 months is negative then the woman can return to routine 5 yearly screening
- If the second repeat HPV test in 12 months is positive (regardless of type) then referral to colposcopy is advised

NB: Women who are overdue for their CST, are over 50 or of ATSI origin, may be referred to colposcopy after the second (rather than third) positive HPV.

3. Higher Risk:

- If the test is positive for HPV 16/18 then referral for colposcopy is advised (regardless of the result of the reflex cytology)
- If the test is positive for other oncogenic HPV types (i.e. NOT 16/18) and the reflex cytology performed by the lab shows possible or definite high grade changes then referral for colposcopy is advised

Please see flowchart* of pathways below:



For women aged 70-74

- If the routine screening HPV test is negative, then the patient is discharged from the screening program and no further screening is required.
- Any positive oncogenic HPV result (regardless of type) should be referred for colposcopy in this age group.
- Women aged 75 years or older who have never had a cervical screening test, or not had one in previous five years may request a test and be screened.

Test of cure for women with treated high-grade intraepithelial lesion (HSIL)

- HPV testing and cytology co-testing should be performed at 12 and 24 months post treatment.
- Once a woman has tested negative by both tests on two consecutive occasions, she is regarded as passing the Test of Cure and can safely return to the normal five-yearly screening interval.
- Any positive test for HPV 16/18 or a cytology result of possible HSIL or HSIL should be referred back to colposcopy.

Follow-up of women treated for Endocervical Adenocarcinoma in-situ (AIS)

- The follow-up of women treated for AIS is annual HPV and cytology co-test indefinitely.
- Any abnormal result will require referral back to colposcopy.
- Currently, the clinical evidence does not support a Test of Cure for endocervical lesions as there is for squamous lesions.
- It is important to inform the pathology laboratory if a woman has a history of AIS so that the appropriate cytology co-test is performed and the appropriate clinical recommendation can be made.

Screening after hysterectomy

- If the hysterectomy was for benign reasons (e.g. fibroids) and there was no history of cervical abnormality, then no follow-up is required.
- If the hysterectomy was for benign reasons (e.g. fibroids) and the cervical screening history is not available, then two consecutive negative HPV tests, 12 months apart, are advised before no further testing is required.
- Any woman with a hysterectomy and a history of HSIL is advised to pass the Test of Cure (as described previously). The Test of Cure can be done either prior to or after the hysterectomy. Once the Test of Cure is passed, then no further follow-up is required.
- Any woman with a hysterectomy and a history of Endocervical AIS should have annual HPV and cytology co-testing indefinitely as described in the previous section.
- Women who have undergone a subtotal hysterectomy (the cervix is not removed) should be invited for 5-yearly HPV testing in accordance with the recommendation for the general population.

Immune-deficient women

- Women with Human immunodeficiency virus (HIV) or a solid organ transplant should have 3-yearly HPV Screening tests.
- Immune-deficient women with any positive oncogenic HPV result (regardless of type) should be referred for colposcopy.
- It is important to inform the pathology laboratory if a patient is immune-deficient so that the appropriate clinical recommendation can be made.

Screening in diethylstilbestrol (DES) exposed women

- Women exposed to DES in utero should be offered an annual HPV and cytology co-test indefinitely.
- Any abnormal result will require referral back to colposcopy.
- Self-collection is not recommended.
- It is important to inform the pathology laboratory if a woman has a history of DES exposure so that the appropriate cytology co-test is performed and the appropriate clinical recommendation can be made.

Investigation of abnormal vaginal bleeding

- Women with abnormal vaginal bleeding should be offered an HPV and cytology co-test.
- Regardless of the test results, referral to gynaecological assessment for investigation of the bleeding should be considered.
- It is important to inform the pathology laboratory if a woman has clinical symptoms so that the appropriate cytology co-test is performed and the appropriate clinical recommendation can be made.

For women under 25 years old

- Routine cervical screening is NOT recommended for asymptomatic women under the age of 25 years.
- For women who experience first sexual activity at a young age (<14 years) and had not received the HPV vaccine before becoming sexually active, a single HPV test between age 20 and 24 can be considered on an individual basis.

Specimen Collection for Cervical Screening

The collection of cervical cells is completed the same way a usual Pap Smear was collected, with the only difference being that the cervical cells are now placed in a ThinPrep® vial, rather than on a slide.

1. Obtain an adequate sample from the cervix

- a) Use luke warm water to lubricate and warm the speculum. A water soluble gel lubricant can be sparingly applied to the posterior blade if necessary. Do not use carbomer-based lubricants.
- b) Insert the speculum.
- c) Insert the central bristles of the Cervical Brush deep enough into the endocervical canal to ensure the shorter bristles contact the exocervix, then push gently and rotate the broom in a full clockwise direction **4 - 5 times**.

Please note: An endocervical Brush should not be used in pregnant women.

2. Place the Cervical Brush into the ThinPrep® vial ASAP

- a) Ensure the ThinPrep® vial is within the use by date. It can be stored at room temperature.
- b) Push the Cervical Brush into the bottom of the vial **10 times**, ensuring to push hard enough to force the bristles apart.
- c) Swirl the Cervical Brush before removing it from the vial.
- d) Discard the Cervical Brush.

Please note: Do not make any glass slides.

3. Secure the cap on the vial

Tighten the cap enough that the torque line on the cap is in line with the torque line on the vial.

4. Record the patient details and complete the request form

- a) Record the patient's full name and date of birth on the vial.
- b) Complete the patient's details on the request form providing as much information as possible.

Pertinent clinical details are essential for reliable cervical screening.

Please ensure to request **Cervical Screening Test, Routine** in screening cases and **Cervical Co-test, Symptomatic** for symptomatic cases.

Other important information should be noted on the request form under clinical notes.

Doctor please consider:

- Patients who are symptomatic (e.g. history of abnormal vaginal bleeding) require both an HPV and a concurrent liquid-based cytology (LBC).
- Patients who have previously been diagnosed with endocervical adenocarcinoma in-situ (AIS), also require a concurrent LBC annually.
- Patients who are immune deficient are advised to repeat testing in 3 years not 5 years.

5. Package the sample and request form for transport

Place both the vial and request form into a specimen bag for transport to the laboratory in the usual manner.

Self-collection

As of July 2022, many more patients are now eligible for self collect. Please refer to 1459_CervScrnSelfCollectGuide for further information.

Turnaround Time

Results will be released to the requesting doctor approximately 7-10 days after the sample is received by the laboratory.

Cost

Cervical screening requests that follow the national prescribed clinical guidelines will be bulk billed subject to Medicare guidelines and criteria.

If Medicare guidelines and criteria are not met, an out-of-pocket fee may apply. If the patient wants additional cervical cytology smears that do not fit the MBS criteria, these tests will not be rebated by Medicare.

Further Information

Brisbane

Dr Rachel Maywald	Head of Cytopathology	(07) 3121 4961
Dr Stephen Peyton	Consultant Cytopathologist	(07) 3121 4961
Dr Ekta Bhutada	Consultant Cytopathologist	(07) 3121 4961

Sunshine Coast

Dr Ethan Oost	Consultant Cytopathologist	(07) 5441 0200
---------------	----------------------------	----------------

Gold Coast

Dr Darina Vuong	Consultant Cytopathologist	(07) 5668 4444
-----------------	----------------------------	----------------

*Source: National Cervical Screening Program: Guidelines for the management of screen-detected abnormalities, screening in specific populations and investigation of abnormal vaginal bleeding. CCA 2016. Accessible from http://wiki.cancer.org.au/australia/Guidelines:Cervical_cancer/Screening. Updated Dec 2020.